Birth Control Options

Please review available contraceptive options and mark wh	nat you would like to discuss further.
☐ Bilateral Tubal Ligation	
 permanent sterilization-performed in the oper 	rating room
* Paragard IUD-intrauterine device-lasts 10 years	G
hormone free	
 Cycles may be heavier and crampier 	
* Mirena IUD-intrauterine device-lasts 5 years	
 Irregular bleeding for the first 6 weeks, may la 	ast 3-6 months. By the end of the first year
cycles may be absent or very light	
* Skyla IUD-intrauterine device-lasts 3 years	
 May have irregular spotting or bleeding at first 	st. Cycles are usually lighter than normal
but may be irregular	, , , ,
□ * Nexplanon-implanted in the upper inner arm-lasts	3 years
 May have irregular bleeding that does not res 	•
☐ Combined Oral Contraception-pills with estrogen	, ,
Pill must be taken at a similar time daily	
 Not usually prescribed while breastfeeding 	
☐ Nuva Ring	
 Vaginal ring inserted once a month and remo 	oved for a monthly cycle
 Not usually prescribed while breastfeeding 	
□ Progesterone Only Pill	
 Pill must be taken at the same time daily 	
 May cause more irregular bleeding than the or 	combined oral contraceptive
☐ Condoms	
 Must be used correctly to avoid pregnancy 	
□ Diaphragm	
 Device is fitted in the office 	
 Must be used correctly to avoid pregnancy 	
Natural Family Planning	
 May be unpredictable when breastfeeding 	
Rely on significant other's vasectomy	
* A device with an * by it indicates that it needs to be ordered. Ye	ou will need to sign for the device and it may
take up to 6 weeks to receive this device. Verification of benefits	·
obtained through your insurance company prior to ordering. Pay	•
will be due at the time of service. You should avoid sex for two v	
provider may elect to reschedule your appointment if there is con	•
Name printed	Date of birth
Signature	Date